

## 心研データベースを用いた最近の研究実績

### < 2008 年 >

1) Suzuki S, Yamashita T, Ohtsuka T, Sagara K, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Ogasawara K, Sawada H, Aizawa T.  
Prevalence and Prognosis of Patients With Atrial Fibrillation in Japan. *Circ J.* 2008;72:914-920.

### < 2009 年 >

2) Suzuki S, Yamashita T, Otsuka T, Sagara K, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Ogasawara K, Sawada H, Yamazaki T, Aizawa T.  
Treatment strategy and clinical outcome in Japanese patients with atrial fibrillation. *Heart Vessels.* 2009;24:287-93.

3) Nakamura M, Yamashita T, Yajima J, Oikawa Y, Ogasawara K, Kirigaya H, Sagara K, Koike A, Sawada H, Aizawa T.  
Impact of reduced renal function on prognosis in Japanese patients with coronary artery disease: a prospective cohort of Shinken Database 2007. *Hypertens Res.* 2009;32:920-6.

4) Nakamura M, Yamashita T, Yajima J, Oikawa Y, Ogasawara K, Kirigaya H, Nagashima K, Sawada H, Aizawa T.  
Long-term safety and efficacy of sirolimus-eluting stents in Japanese patients: a single-center cohort study. *J Invasive Cardiol.* 2009;21:526-31.

5) Goda A, Yamashita T, Suzuki S, Ohtsuka T, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Sagara K, Ogasawara K, Isobe M, Sawada H, Aizawa T.  
Prevalence and prognosis of patients with heart failure in Tokyo. *Int Heart J.* 2009;50:609-25.

### < 2010 年 >

6) Suzuki S, Yamashita T, Ohtsuka T, Sagara K, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Ogasawara K, Sawada H, Yamazaki T, Aizawa T.  
Body Size and Atrial Fibrillation in Japanese Outpatients. *Circ J.* 2010;74:66-70.

7) Nakamura M, Yamashita T, Yajima J, Oikawa Y, Ogasawara K, Sagara K, Kirigaya H, Koike A, Nagashima K, Ohtsuka T, Uejima T, Suzuki S, Sawada H, Aizawa T.  
Clinical outcome after acute coronary syndrome in Japanese patients: an observational cohort study. *J Cardiol* 2010;55:69-76

8) Nakamura M,  
Brachial-ankle Pulse Wave Velocity as Risk Stratification Index of Short-term Prognosis in Type 2 Diabetic Patients with Coronary Artery Disease. *Hypertens Res.* 2010;33:1018-24

9) Goda A, Yamashita T, Suzuki S, Ohtsuka T, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Sagara K, Ogasawara K, Isobe M, Sawada T, Aizawa T.  
Heart failure with preserved versus reduced ventricular systolic function: a prospective cohort of Shinken Database 2004-2005: *J Cardiol* 2010;55:108-116

< 2011 年 >

10) Suzuki S, Yamashita T, Otsuka T, Sagara K, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Ogasawara K, Sawada H, Aizawa T.

Recent mortality of Japanese patients with atrial fibrillation in an urban city of Tokyo. *J Cardiol* 2011;58:116-123.

11) Nakamura M, Yamashita T, Yajima J, Oikawa Y, Ogasawara K, Sagara K, Koike A, Kirigaya H, Nagashima K, Otsuka T, Uejima T, Funada R, Matsuno S, Suzuki S, Sawada H, Aizawa T.

Shinken Database Study Group. Impact of early statin initiation on secondary prevention in Japanese patients with coronary artery disease. *J Cardiol* 2011;57:172-180.

< 2012 年 >

12) Suzuki S, Sagara K, Otsuka T, Matsuno S, Funada R, Uejima T, Oikawa Y, Koike A, Nagashima K, Kirigaya H, Yajima J, Sawada H, Aizawa T, Yamashita T.

Gender-Specific Relationship Between Serum Uric Acid Level and Atrial Fibrillation Prevalence. *Circ J* 2012;76:607-611.

13) Suzuki S, Otsuka T, Sagara K, Matsuno S, Funada R, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Sawada H, Aizawa T, Yamashita T.

Dabigatran in Clinical Practice for Atrial Fibrillation With Special Reference to Activated Partial Thromboplastin Time. *Circ J* 2012;76:755-757.

14) Suzuki S, Sagara K, Otsuka T, Matsuno S, Funada R, Uejima T, Oikawa Y, Koike A, Nagashima K, Kirigaya H, Yajima J, Sawada H, Aizawa T, Yamashita T.

A New Scoring System for Evaluating the Risk of Heart Failure Events in Japanese Patients with Atrial Fibrillation. *Am J Cardiol*;110:678-682.

15) Seno K, Suzuki S, Sagara K, Otsuka T, Matsuno S, Funada R, Uejima T, Oikawa Y, Yajima J, Koike A, Nagashima K, Kirigaya H, Sawada H, Aizawa T, Yamashita T.

Distribution of first-detected atrial fibrillation patients without structural heart diseases in symptom classifications. *Circ J* 2012;76:1020-1023.

16) Kaneko H, Koike A, Seno K, Tanaka S, Suzuki S, Nagayama O, Sagara K, Otsuka T, Matsuno S, Funada R, Uejima T, Oikawa Y, Yajima J, Nagashima K, Kirigaya H, Sawada H, Aizawa T, Yamashita T.

Role of cardiopulmonary dysfunction and left atrial remodeling in development of acute decompensated heart failure in chronic heart failure with preserved left ventricular ejection fraction. *J Cardiol*;59:359-365.

< 2013 年 >

17) Suzuki S, Sagara K, Otsuka T, Kano H, Matsuno S, Uejima T, Oikawa Y, Koike A, Nagashima K, Kirigaya H, Yajima J, Sawada H, Aizawa T, Yamashita T.

Estimated glomerular filtration rate and proteinuria are associated with persistent form of atrial fibrillation: Analysis in Japanese patients. *J Cardiol* 2013;61:53-57.

18) Suzuki S, Sagara K, Otsuka T, Kano H, Matsuno S, Takai H, Uejima T, Oikawa Y, Koike A, Nagashima K, Kirigaya H, Yajima J, Tanabe H, Sawada H, Aizawa T, Yamashita T.

Usefulness of Frequent Supraventricular Extrasystoles and a High CHADS2 Score to Predict First-Time Appearance of Atrial Fibrillation. *Am J Cardiol* 2013;111:1602-1607.

19) Kaneko H, Yajima J, Oikawa Y, Tanaka S, Fukamachi D, Suzuki S, Sagara K, Otsuka T, Matsuno S, Funada R, Kano H, Uejima T, Koike A, Nagashima K, Kirigaya H, Sawada H, Aizawa T, Yamashita T.

Impact of aging on the clinical outcomes of Japanese patients with coronary artery disease after percutaneous coronary intervention. *Heart Vessels*; in press.

20) Kaneko H, Yajima J, Oikawa Y, Tanaka S, Fukamachi D, Suzuki S, Sagara K, Otsuka T, Matsuno S, Funada R, Kano H, Uejima T, Koike A, Nagashima K, Kirigaya H, Sawada H, Aizawa T, Yamashita T.

Effects of statin treatment in patients with coronary artery disease and chronic kidney disease. *Heart Vessels*; in press.